



GROWTH COMPONENTS, FORAGE AND SEED YIELD OF VELVET BEAN (*Mucuna pruriens*) AS AFFECTED BY PIG MANURE RATES AND IRRIGATION INTERVALS IN NIGERIAN SAVANNA

¹Hassan, M. R., ¹Abdu, S.B., ²Amodu, J. T., ¹Uchechukwu, A. V., ¹Hasiya, I., ¹Adamu, H.Y., ¹Musa, A., ¹Braimah, Y., ³Sani, I., ⁴Akpensuen, T. T., ²Ishiaku, Y. M., ⁵Bala, A. G. and ¹Bello, S. S.

¹Department of Animal Science, Ahmadu Bello University, Zaria.

²Feeds and Nutrition Research Programme, National Animal Production Research Institute, Ahmadu Bello University, Zaria.

³Department of Animal Production, Faculty of Agriculture and Agricultural Technology, Abubakar Tafawa Balewa University, Bauchi, Bauchi State, Nigeria. ⁴Department of Animal Science, Faculty of Agriculture, University of Jos, Plateau State.

⁵Division of Agricultural Colleges, College of Agriculture and Animal Science, Mando, Kaduna

*Corresponding author: mrhassan@abu.edu.ng Tel: +23480628533

Abstract

A study was conducted to determine the effects of graded levels of pig manure rates and irrigation frequency on growth components, forage and seed yield of Velvet bean (*Mucuna pruriens*). The experimental design was a split plot fitted into a Randomized Complete Block Design (RCBD), replicated three times. Treatments consisted of graded levels of pig manure application rates (0, 9 and 18 t / ha) and three irrigation frequencies (2,4 and 6 days intervals). Results indicated that tallest plant, (112.44cm) (P<0.05), more leaves (107.75) were obtained when pig manure was applied at 18 t / ha and irrigated at 6 days irrigation interval, which was at par with result applied 9 t / ha at 4 days irrigation interval. However, the leaf area index (LAI) did not differ (P>0.05) with increase in irrigation interval. Also, the heaviest (P<0.05) fresh and dry forage yields (9.22t/ha vs. 6.66t/ha) were observed when pig manure was applied at 18 t / ha, at irrigation interval of 6 days, which was at par with applied 9 t/ha and 2 days irrigation interval. For seed production, irrigation interval of 2 days produced the highest (P<0.05) yield (2.98t/ha), when pig manure was applied at the rate of 9 t / ha. It was therefore concluded that pig manure could be applied to *Mucuna pruriens* forage at 9 t / ha at irrigation interval of 4 days, for better growth components. However, for forage yield and seed production, 9t/ha of pig manure and 2 days irrigation interval produced the best result in Northern Guinea savannah of Nigeria.

Key words: Dry matter, irrigation, Nigeria, nutrients, soil, velvet bean.

INTRODUCTION

Mucuna (*Mucuna pruriens* L.), is a popular leguminous forage that is used for different purposes in tropical countries (Kevin *et al.*, 2008; Anastasia and Joseph, 2003). It

belongs to the family Fabaceae (*leguminoceae*) and subfamily *Faboideae* (USFS, 2011). There are many species of *Mucuna* which are characterized by remarkable physical and physiological

variations (Pengelly *et al.*, 2004). The black seeded *Mucuna pruriens* has superior quality than others, especially in early seed ripening and vegetative growth (Onoma and Asongwed, 2002). *Mucuna* is believed to have originated from Eastern India, Malaysia, and southern parts of China (Janardhanan and Lakshaman, 1985; FAO, 2011).

Mucuna is cherished by farmers as a leguminous forage crop used for livestock feeding and green manuring in different parts of the tropics (Mbutia and Gachuri, 2003) that is compatible with other pasture species (Whitbread *et al.*, 2004). It also has the ability to smother weeds with less problem of insect attack (FAO, 2011, Agyenim, 2006). Many cultivars of *Mucuna* are suitable for humid regions of the tropics, while others are suitable for dry land farming (Anastasia and Joseph, 2003). *Mucuna* can be grown successfully on soils unsuitable for cowpeas, and therefore helps to protect the soil from environmental hazards. However, it has the disadvantage of a longer growth period and the seeds are more difficult to thresh. Moreover, the seeds are not highly valued for human consumption or animal feeding, because of the prolonged soaking and, or boiling required before it can be cooked for safety consumption (Ross, 1999). The leaves and vines make an excellent fodder for livestock (Westphal, 1974).

In developing countries such as Nigeria, *Mucuna* is produced by smallholder farmers for soil fertility improvement and livestock production (Ngosong *et al.*, 2015). However, the major challenges include drought (Gabrielle, 2016), scarcity and cost of initial fertilizer doses to be applied for proper vegetative growth and seed yield (Odedina *et*

al., 2011). The use of animal manure such as cattle, sheep, poultry and pig manure in crop production with the aim of reducing production costs and environmental damages has been reported (Odedina *et al.*, 2011; Obi and Ebo, 1995; Ojeniyi, 2002). Despite its ability to grow on soils with low available soil phosphorus, *Mucuna* has been reported to respond to phosphorus and lime applications on acid soils, either from amelioration of pH or from Mg and Ca applications (Maasdorp *et al.*, 2004). The use of *Mucuna* seeds to supplement poor quality grasses in small ruminant diet has been reported in Zimbabwe (Matenga *et al.*, 2003). Pig manure is a very good source of organic matter which plays a vital role in improving the physical, chemical and biological properties of the soil, which activate the activities of soil microorganism, and enhance slow release of plant nutrients (macro and micro) with resultant environmental harmony (Sharma and Mitta, 1991). Efficient and effective use of pig manure ensure sustainable crop productivity, because soil organic matter is maintained, nutrients are released more slowly, stored for a longer time in the soil, improved root development and high crop yield (Sharma and Mitta, 1991).

Global climate change (GCC) has led to an increasing instability differences in the rainfall distribution pattern for crop cultivation. This leads to developing methods of irrigation that will minimize the impact of drought, thereby maximizing the crop water use efficiency (Salah *et al.*, 1995). Irrigation is an important component of the farming system in Nigeria. It helps to supply enough moisture to crops for better performance. Similarly, irrigation fetches

more income to farmers than rainfed farming. In Nigeria, smallholder farmers hardly utilize pig manure as a source of fertilizer for their crops, despite its availability and low cost. Therefore, it was hypothesized that utilization of pig manure by smallholder farmers, with appropriate agronomic practices, will help to provide cheap and sustainable option for forage crop production under irrigation in Nigeria. This will lead to improvement in productivity of livestock and economic status of farmers thereby preventing further environmental degradation, due to indiscriminate use of chemical fertilizers by the farmers. Hence, this study was conducted to evaluate the effect of single application of graded levels of pig manure on growth components, forage and seed yield of *Mucuna pruriens* under different irrigation frequencies in Northern Guinea savannah of Nigeria.

MATERIALS AND METHOD

The study was carried out at Samaru located between latitude 11⁰11'N and longitude 7⁰38'E, on altitude of 686 m above sea level. The area falls within the Northern Guinea Savannah zone with an average rainfall of 1100mm. The maximum temperature varies from 26⁰C to 35⁰C depending on the season, while the mean relative humidity during harmattan period and the wet season are 21% and 72%, respectively (Ovimaps, 2016). Seeds of *Mucuna pruriens* used for the study were obtained from the Feeds and Nutrition Research Programme at the National Animal Production Research Institute (NAPRI) Shika, Zaria. The soil at the experimental site was taken prior to land preparation using soil auger at 0-15cm depth for routine analysis, which was conducted in

the Department of Soil Science, Faculty of Agriculture, Ahmadu Bello University, Zaria. Table 1 shows the physical and chemical properties of the soil at the experimental site.

Treatments, Experimental Design, and Data Collection

The treatments consist of graded levels of pig manure application (0, 9 and 18 t/ha) and three irrigation intervals (2, 4 and 6 days intervals). The experimental design was a split plot fitted into a Randomized Complete Block Design (RCBD), replicated three times. The land was cleared, prepared and irrigated for 7 days during the 2014/2015 dry season to attain field capacity. Net plots measuring 1.5m by 1.5m were prepared using hoes. Seeds of *Mucuna* (black) were treated with Apron star, which is a fungicide at the rate of 4 kg/ sachet and then sown using dibbling method at inter and intra row spacing of 25cm, respectively. All plots were fertilized with a basal NPK fertilizer at the rate of 60kg/ha.

Data were collected for agronomic indices at 12 weeks after sowing on plant height, number of leaves, leaf length, leaf breadth, number of branches, number of nodes and fresh forage yield/ha. Plant height, leaf length and breadth were measured using a meter rule. Fresh forage yield was estimated using a 0.5 m by 0.5 m quadrant in each plot. The forage material was cut using a hand sickle at 10cm height, bulked and weighed immediately using a hanging scale before sub-sampling 300g per plot for the determination of proximate composition of the forage. Forage samples were oven dried at 60°C for 72 hours until they attained constant weight. The dried samples were

milled to pass 1mm mesh using an electric miller. Milled samples of *Mucuna pruriens* forage were analyzed for proximate composition: dry matter(DM), crude fibre(CF), crude protein(CP% =N×6.25), nitrogen free extract (NFE), ether extract(EE) and ash content, respectively at the Biochemistry Laboratory of the Department of Animal Science, Ahmadu Bello University, Zaria, using the method described by A O A C. (2005). All data collected were subjected to analysis of variance (ANOVA) using ‘F’ test as the Statistical Analysis System (2002) computer software package. Means were compared by a Post-hoc Tukey’s HSD test ($P<0.05$). Since there were significant interactions, only main effects of factors were considered in the results.

RESULTS AND DISCUSSION

The result of the soil analysis shows that the soil was a sandy-loam under the soil textural classification (Table 1).

The pH was 5.60 which indicated that the soil was slightly acidic. The result shows that the total nitrogen and organic carbon contents of the experimental site were low. This showed that there was need for the application of nutrients to support full establishment and seed production of *Mucuna pruriens*. The chemical composition of pig manure showed that the manure was rich in essential nutrients (Table 2).

Many studies reported positive improvements in growth performance of different crops fertilized with pig manure (Ojeniyi and Akanni, 2008; Olomilua *et al.*, 2007). Findings of Sanni and Okeowo (2016) confirmed the superiority of using pig manure on eggplant compared to other

manure sources. Table 3 shows the results of interaction effect of levels of pig manure application and irrigation frequency.

The tallest plants (112cm) and more of leaves per plant (108) were observed at combination levels of 18 t / ha pig manure rate and 4 days irrigation interval and 18 t / ha and 6 days irrigation interval, respectively. This result is similar to the findings of Sanni and Okeowo (2016) in eggplant. These authors found that plots treated with pig manure at the rate of 7.5 t / ha produced the highest mean height of eggplant. Results obtained in this study might be related to the high nutrient content of pig manure, especially calcium and phosphorus, which are necessary for plant growth (Odedina *et al.*, 2011; Adeniyi and Ojeniyi, 2003). It was also reported that pig manure could increase the contents of soil macro and micro nutrients and yield of different crops (Awodun and Alafusi, 2007; Olomilua *et al.*, 2007; Ayeni *et al.*, 2009). Results of Okonmah (2011) on okra plants showed that plants that received pig manure at the rate of 15t/ha, produced the best growth parameters, which is supported by the findings of Giwa and Ojeniyi (2004) and Christo *et al.* (2011), proving the efficiency of pig manure in supporting plant growth. The highest number of leaves of *Mucuna* recorded in this study (108) obtained at treatment combination of 18 t / ha pig manure rate and 6 days irrigation interval was significantly lower than the value reported in eggplant (239) at lower rate of pig manure application (7.5t/ha). This result can be explained by the excessive leaf shading effect of *Mucuna* forage at maturity period, due to rapid formation of new dark green leaves (Zsolany and Gorlitz, 1994).

Similarly, the highest quantity of pig manure applied in this study might have led to higher accumulation of nitrogen in the soil. However, the rate of nitrogen uptake from the soil by *Mucuna* forage is minimum compared to non-nitrogen fixing plants such as okro plant, since it has been reported that the rate of nitrogen uptake by plants is proportional to the quantity of nitrogen in the soil (Madukwe *et al.*, 2008). Irrigation interval of 4 or 6 days in combination with pig manure rate of 18 t / ha showed the tallest plants and number of leaves of *Mucuna* forage, respectively (Table 3). This indicated that *Mucuna* forage could survive dry spell condition, which further shows that it can be successfully produced in areas with less rainfall in Nigeria. Salah *et al.* (1995) reported a similar effect of irrigation interval on agronomic and quality indices of alfalfa forage in Riyadh, Saudi Arabia. The leaf area index of *Mucuna* forage in this study was however statistically similar across the treatments. This result might be related to the rapid canopy development and short photoperiod in legumes. Lawson *et al.* (2013) reported reduced leaf area index in cowpea intercropped with cereal in Ghana. However, the result obtained in this study is contrary to the findings of Nweke *et al.* (2008) who reported an effect of different nutrient sources on the leaf area index of maize crop. These authors showed that combination of pig manure and urea fertilizer produced the best leaf area index. Table 4 shows the results of combined effect of pig manure application and irrigation frequency on forage yield of *Mucuna pruriens*.

Results obtained indicated that the highest fresh (9.22t/ha) and dry forage yields (6.66t/ha) were recorded in all applied

manure rates at 6 days irrigation interval, which was also at par with applied 9 t / ha of pig manure at 2 days interval, except on dry forage yield where depression was observed with applied 9 t / ha at 6 days irrigation interval. This suggests that *Mucuna* forage can also survive short term water logging, in soils with low nutrient status in Northern Guinea savannah of Nigeria. The values obtained in this study fall within the range of 5-12t/ha previously reported (Maasdorp *et al.*, 2004). The *Mucuna* forage in this study was cut once at maturity stage of growth (16 weeks), which could explain the reason for the general low forage yield obtained in this study. Ross (1999) reported that when *Mucuna* plant is intended for forage, it can be harvested at the period of 90-120 days. Increasing the irrigation interval from 2 to 6 days in this study led to progressive increase in both fresh and dry forage yields. This result could be related to the fact that *Mucuna* plant has the ability to tap adequate soil moisture through its long tap root system in addition to unintercepted sunlight received. However, Salah *et al.* (1995) reported dramatic decreases in forage yields of alfalfa in Riyadh in fresh (97.3 vs. 91.7t/ha/yr) and dry forms (25.3 vs 23.1t/ha/yr), when irrigation interval was increased from 5 to 10 days interval, respectively. Donovan and Meek (Donovan and Meek, 1983), reported that proper irrigation of legumes such as alfalfa is a critical factor in the success of the crop. Also, research has shown that water stress condition has a remarkable decrease in top and root growth of legumes and grass species (Stevenson and Laidlaw, 1985). Maasdorp *et al.* (2004) suggested that

growth of *Mucuna* is affected by continuous moisture supply in the soil, especially when they are defoliated before flowering.

Table 5 presents the result of seed yield of *Mucuna pruriens* as influenced by level of pig manure application and irrigation interval. The highest ($P<0.05$) seed yield (2.89t/ha) was produced at 9 t / ha of pig manure application at 2 days irrigation interval. However, increasing the level of pig manure application rate and irrigation interval to 18t/ha and 4 days irrigation interval, led to 13% decrease ($P<0.05$) in the quantity of seeds produced by *Mucuna*. This indicates that for seed production, farmers in the study area should irrigate *Mucuna* forage at 2 days interval with 9t/ha of pig manure as a source of nutrients, for proper flowering and seed production. Although the results obtained showed that both seed yields at 9 and 18 t / ha at irrigation intervals of 2 and 4 days are at par, but using lower rate of pig manure might be more economical to farmers in Nigeria. Similar findings were reported by Eilittä and Carsky (2003), Omokanye (2001) and Lal (2015).

This supports the fact that higher rates of pig manure application on crops might results in highest seed yield, although Ojeniyi and Olamilua (2006) observed the highest yield of okro pods at 10t/ha pig manure application rate. Work by Christo *et al.* (2011) on fruit production by eggplant supported that pig manure can serve as good substitute to poultry manure.

CONCLUSION

The results of this study showed that for production of fresh forage and dry matter yields as well as seed yield of *Mucunapruriens*, farmers in the study area

should apply pig manure at the rate of 9t/ha and irrigate at 2 days interval for better yield.

REFERENCES

- A.O. A. C. (2005): Official Methods of Analysis, 15th edition. Association of Official Analytical Chemists, Washington, D.C.
- Adeniyani, O.N. and Ojeniyi, S. O. (2003). Comparative effectiveness of different levels of poultry manure with NPK fertilizer on residual soil fertility nutrient uptake and yield of maize. *Moor Journal of Agricultural Resources*, 4, 191-197.
- Agyenim Boateng, S. (2006). *Mucuna pruriens* and its Effect on some Physical, Chemical and Biological Properties of a Forest Acrisol, *West African Journal of Application Ecology*, 35 (8, 1): 393.
- Anastasia, A, A. and Joseph, O. (2003). Variability in productivity of *Mucuna pruriens* varieties in a semi-arid environment. Jean-Yves Jamin, Lamine Seiny Boukar, Christian Floret. *Cirad - Prasac*, 6 p. <hal-00137962>
- Awodun, M.A. and Alafusi, R.A. (2007): Effect of integrated application of goat manure and urea on soil nutrient content and yield of Okra. *Nigerian Journal of Soil Science*, 17, 126-129.
- Ayeni, L.S., Adetuniji, M.T. and Ojeniyi, S.O. (2009). Integrated application of NPK fertilizer, Cocoa pod ash and poultry manure: effect on maize performance, plant and soil nutrient content. *International Journal of*

Growth Components, Forage and Seed Yield of Velvet Bean (*Mucuna pruriens*) As Affected by Pig Manure Rates and Irrigation Intervals in Nigerian Savanna

- Pure and Application Science*, 2(2), 34-41.
- Christo, I. E., Okorie, H. A. and Chikere, N. C. (2011). Performance of Eggplant (*Solanum gilo* L.) as affected by manure types and rates, *Global Resource. Journal of Science*, 1:43-47.
- Donovan, T. J. and Meek, B. D. (1983). Alfalfa Responses to Irrigation Treatment and Environment. *Agronomy Journal*, 75:461-464.
- Eilittä, M., Carsky, R. J. (2003). Efforts to improve the potential of *Mucuna* as a food and feed crop: background to the workshop. *Tropical and Subtropical Agro-ecosystem*, 1: 47-55.
- FAO, 2011. Grassland Index. A searchable catalogue of grass and forage legumes. FAO, Rome, Italy.
- Gabrielle, K. (2016). Pulse crops and sustainability: A framework to evaluate multiple benefits. Lexeme Consulting. *Pulses*, 1-64.
- Giwa, D. D. and Ojeniyi, S.O. (2004) Effect of integrated application of pig manure and NPK on soil nutrient content and yield of tomato (*Lycopersicon esculentum* Mill.) Proceedings of the 29th Annual Conference of the Soil Science Society of Nigeria, pp 164-169.
- Janardhanan, K.V. and Lakshaman, A.R. (1985). Studies on chemical composition and anti nutritional factors in three germplasm seed material of the tribal pulse *Mucuna pruriens* www.sciencedirect.com/science/article/pii/0308814692902357
- Kevin, L. A., Kenneth, A. A., Joseph, G. L. and Heathcliffe, R. (2008). Intercropping Corn with Lablab Bean, Velvet Bean, and Scarlet Runner Bean for Forage. *Crop Science*, 48:371–379.
- Lal, R.K. (2015). Genotype selection for agronomical trial: Seed yield in Kewachh {*Mucuna pruriens* (L)}. *Industrial Crops and Products*, 65: 62-70.
- Lawson, I.Y.D., Issahaku, A., Acheampong, S.K., Adams. B. And Tuffour, V. (2013). Time of planting and weed suppression abilities of some legumes intercropped with maize in the Guinea savanna zone of Ghana. *Agricultural and Biology Journal of North America*, 4(4): 358-363.
- Maasdorp, B., Jiri, O. and Temba, E. (2004) Contrasting adoption, management, productivity and utilization of *Mucuna* in two different smallholder farming systems in Zimbabwe. In: Matenga, V.R., Ngongoni, N.T., Titterton, M., Maasdorp, B.V.(2003). *Mucuna* seed as a feed ingredient for small ruminants and effect of ensiling on its nutritive value. *Tropical and Subtropical Agro-ecosystem*, 1 (2-3): 97-105.
- Madukwe, D.K., Christo, I. E. C. and Onuh, M. O. (2008). Effect of Organic Manure and Cowpea Varieties (*Vigna unguiculata* (L) Walp), varieties on the chemical properties of soil and root nodulation. *Science World Journal*, 3(1): 43-46.

- Mbuthia, E.W. and Gachuri, C.K. (2003). Effect of inclusion of *Mucuna pruriens* and *Dolichos lablab* forage in Napier grass silage on silage quality and on voluntary intake and digestibility in Sheep. *Tropical and Subtropical Agro-ecosystem*, (1): 123 – 128.
- Ngosong, C., Mfombep, P. M., Njume, A. C. and Tening, A. S. (2015). Integrated soil fertility management: Impact of *Mucuna* and *Tithonia* biomass on Tomato (*Lycopersicon esculentum*) performance in smallholder farming system. *Agricultural Sciences*, 6:1176-1186.
- Nweke, I.A. and Obasi, M.N. (2008). Effect of Different levels of pig manure on the growth and yield of Okra (*Abelmoschus esculentum* L. Moench). Proceedings of Agricultural Society of Nigeria, 47th Annual conference, held at Ibadan: 23-25pp.
- Obi, M.E. and Ebo, P.O. (1995). The effect of organic and inorganic amendments on soil physical properties and maize production in severely degraded soils in southern Nigeria. *Biological Resource and Technology*, 51:117-123.
- Odedina, J. N., Odedina, S. A. and Ojeniyi, S. O. (2011). Effect of Types of Manure on Growth and Yield of Cassava (*Manihot esculenta*, Crantz), *Researcher*, 3(5): 1-8.
- Ojeniyi, S.O. (2002). Effect of goat manure on soil nutrient and okra yield in a rain forest area of Nigeria, *Applied and Tropical Agriculture*. 5:20-23.
- Ojeniyi, S.O. and Akanni, D.I. (2008). Effect of animal manures on soil properties, leaf nutrient composition, growth and yield of pepper (*Capsicum annum*). *Journal of Resource in Agriculture*, 5(4), 86-90.
- Ojeniyi, S.O. and Olamilua, A. I. (2006). Effects of poultry manure on nutrient composition, growth and yield of okra. Proceedings of the 23rd Annual Conference of the Horticulture Society of Nigeria. Pp. 136-139.
- Okonmah, L. U. (2011). The effects of pig manure rates on some agronomic parameters of three okra (*Abelmoschus esculentus* (L) Moench) cultivars in the Asaba Agro-ecological zone. *International Journal of Agricultural Science*, 1 (7):427-436.
- Olomilua, A.I., Akanbi, O.S.O. and Ojeniyi, S.O. (2007): Effect of pig manure on nutrient composition, growth and yield of Okra. *Nigeria Journal of Soil Science*, 17, 109-112.
- Omokanye, O. T. (2001). Seed production of Centro (*Centrosema pubescens*) in the year of establishment at Shika, Nigeria, *Tropiculture*, 19(4):176-179.
- Onoma, J. and Asongwed, A. (2002). Variability in productivity of *Mucuna pruriens* variety in Semi-Arid Environment. Institute of Agricultural Research for Development.
- Ovimaps (2016). Ovi location map; ovi earth imaginary.
- Pengelly, B.C., Whitbread, A., Mazaiwana, P.R., Mukombe, N. (2004). Tropical forage research for the future - Better

Growth Components, Forage and Seed Yield of Velvet Bean (*Mucunapruriens*) As Affected by Pig Manure Rates and Irrigation Intervals in Nigerian Savanna

- use of research resources to deliver adoption and benefits to farmers. In: *Tropical legumes for sustainable farming systems in Southern Africa and Australia*.
- Ross, I.A. (1999). Medicinal Plants of the World. Chemical Constituents, Traditional Uses and Modern Medicinal Uses. Humana Press, Totowa, New Jersey. 415 pp. 1600.
- Salah, S., Din, T. E. and Assaeed, M. A. (1995). Effect of phosphorus fertilizer and irrigation frequency on yield and protein content of alfalfa, *Journal of King Saudi University*, 7: 49-60.
- Sanni, K. O. and Okeowo, T. A. (2016). Growth, yield performance and cost benefit of eggplant (*Solanum melongena*) production using goat and pig manure in Ikorodu, Lagos, Nigeria. *International Journal of Science Resource and Engineering Studies*, 3(4):1-6.
- SAS Institute (2002). SAS/STAT User's Guide version 8.2 edition: Statistics, SAS Institute Incorporation, Cary, New York City.
- Sharma, A.R. and Mitta, B.N. (1991). Effect of different rates of application of organic nitrogen fertilizers in rice based cropping system. *Journal of Agricultural Science, (Cambridge)* 117:313-318.
- Stevenson, C.A. and Laidlaw, A.S. (1985). The effect of moisture stress on stolon and adventitious root development in white clover (*Trifolium repens* L.). *Plant and Soil*, 85:249-257.
- USFS-United States Forest Service (2011). *Mucuna pruriens* (L.) DC. Pacific Island Ecosystems at Risk (PIER).
- Westphal, E. (1974). *Mucuna pruriens* (L) DC. Pulses in Ethiopia, their taxonomy and agricultural significance. pp. 121-129. Wageningen, PUDOC, Cent. Agric. Publ. Doc., *Agricultural Resources*, Rep. 815: 278.
- Whitbread, A.M., Jiri, O. and Maasdorp, B. (2004) The effect of managing improved fallows of *Mucuna pruriens* on maize production and soil carbon and nitrogen dynamics in sub-humid Zimbabwe. *Nutrition Cycling in Agro-ecosystem*, 69, 59–71.
- Zsolany, A. and Gorlitz, G. (1994). Water extractable organic matter in arable soils. Effect of drought and long term fertilization. *Soil Biology and Biochemistry*, 26(9):1257-1261.

Table 1: Physical and chemical properties of the soil at the experimental site

Properties	Value
Physical (%)	
Clay	14
Silt	26
Sand	60
Textural class	Sandy-loam
Chemical:	
pH (H ₂ O)	5.60
pH (CaCl ₂)	5.00
Exchange cations (Cmol/kg):	
Na	0.27
K	0.33
Ca	4.36
Mg	0.84
Total N (%)	0.28
Available P (ppm)	33.25
Organic carbon (%)	0.40
Cation Exchange Capacity (Meq/100g)	8.20
Exchangeable acidity (Cmol/kg):	
H+Al ³⁺	0.40

Table 2: Chemical composition of pig manure applied to *Mucuna pruriens* forage

Parameter	Value
N total (%)	2.66
P (%)	0.23
K (%)	0.11
P/N	0.09
K/N	0.04

Tabl 3: Effect of pig manure application and irrigation frequency on growth components of *m mucuna pruriens*

Irrigation interval (days)	Levels of pig manure applied (t/ha)		
	0	9	18
	Plant height (cm)		
2	84.21 ^a	75.88 ^b	76.67 ^b
4	69.33 ^b	83.33 ^a	112.44 ^a
6	68.47 ^b	60.94 ^c	110.10 ^a
Overall mean (cm)	74.03	73.39	99.92
SEM	2.87	2.86	3.33
LOS	*	*	*
T×I	*	*	*
	Number of leaves (no.)		
2	69.00 ^b	76.88 ^b	80.00 ^c
4	60.38 ^c	85.50 ^a	97.29 ^b
6	76.38 ^a	64.67 ^c	107.75 ^a
Overall mean (no.)	68.58	75.68	95.01
SEM	2.76	2.89	3.25
LOS	*	*	*
T×I	*	*	*
	Leaf Area Index (LAI)		
2	0.15	0.17	0.18
4	0.13	0.16	0.20
6	0.13	0.15	0.21
Overall mean	0.14	0.16	0.19
SEM	0.12	0.13	0.15
LOS	NS	NS	NS
T×I	NS	NS	NS

^{abc} Means with different superscripts along the column differed significantly (P<0.05); LOS = Level of significant; SEM = Standard error of the mean, T×I = Interaction between pig manure application rate and irrigation interval.

Table 4: Effect of pig manure application and irrigation frequency on forage yield of *Mucuna pruriens*.

Irrigation interval (days)	Level of pig manure application (t/ha)		
	0	9	18
	Fresh forage (t/ha)		
2	5.12 ^c	7.74 ^a	8.49 ^b
4	6.33 ^b	6.15 ^b	8.42 ^b
6	7.00 ^a	7.56 ^a	9.22 ^a
Overall mean (t/ha)	6.15	7.15	8.71
SEM	0.11	0.13	0.15
LOS	*	*	*
T×I	*	*	*
	Dry forage yield (t/ha)		
2	2.98 ^c	4.11 ^a	5.38 ^b
4	3.10 ^b	3.77 ^c	5.00 ^c
6	4.57 ^a	3.99 ^b	6.66 ^a
Overall mean (t/ha)	3.55	3.96	5.68
SEM	0.03	0.01	0.02
LOS	*	*	*
T×I	*	*	*

^{abc} Means with different superscripts along the column differed significantly (P<0.05); LOS = Level of significant; SEM = Standard error of the mean, T×I = Interaction between pig manure application rate and irrigation interval.

Table 5: Effect pig manure application and irrigation frequency on seed yield of *Mucuna pruriens*

Irrigation interval (days)	Level of pig manure application (t/ha)		
	0	9	18
	Seed yield (t/ha)		
2	2.04 ^a	2.89 ^a	1.33 ^c
4	1.29 ^b	2.00 ^b	2.55 ^a
6	0.49 ^c	0.36 ^c	1.61 ^b
Overall mean (t/ha)	1.27	1.75	1.83
SEM	0.02	0.04	0.01
LOS	*	*	*
T×I	*	*	*

^{abc} Means with different superscripts along the column differed significantly (P<0.05); LOS = Level of significant; SEM = Standard error of the mean, T×I = Interaction between pig manure application rate and irrigation interval.